

SOME GENERALIZATIONS OF F-CONNECTIONS ON DIFFERENTIABLE MANIFOLDS

DUMITRU, Dan

Faculty of Mathematics-Informatics

Spiru Haret University

dandumitru1984@yahoo.com

Abstract

In this article we generalize the notion of F-connection, where F is an almost product structure on a manifold and we give some characterizations of the connections compatible with those structures.

Key words : (ap)-structure, F-connection

AMS classification: 58A05

1. Introduction

Differentiable manifolds with almost product structures (ap)-structures, together with other important structures were investigated by G. Walker [1], Wilmore [2], and Yano [6]. New properties of these structures were established by Legrand [3] and Hsu [4].

In this paper we generalize the notion of F-connection for a connection ∇ on a manifold, where F is an (ap)-structure, by generalizing the (ap)-structures. We also establish some properties of these connections.

Let M be a differentiable manifold of dimension n and T_pM the tangent space to M in the point p with $p \in M$. We denote with $\Gamma(TM)$ the tangent bundle of M , with $F(M)$ the ring of real functions on M , with $D_s^r(M)$ the $F(M)$ module of (r, s) -tensor fields, with $C(M)$ the set of connections on M , and with $D^1(M)$ the $F(M)$ module of sections of M .

Let F be an almost product (ap)-structure on M , that means F is an $(1, 1)$ -tensor field defined on M satisfying $F^2 = I$, where I is the identity. Setting:

$$\Psi_F(\nabla)_X = \frac{1}{2}(\nabla_X + F \circ \nabla_X \circ F) \text{ and} \tag{1}$$

$$\chi_F(\tau) = \frac{1}{2}(\tau_X + F \circ \tau_X \circ F),$$

where τ is an $(1, 2)$ -tensor field on M , $\forall X \in D^1(M)$ it follows

$$\Psi_F(\nabla) \in C(M), \chi_F(\tau) \in D_2^1(M) \quad (2)$$

and

$$\Psi_F^2 = \Psi_F, \chi_F^2 = \chi_F, \Psi_F(\nabla + \tau) = \Psi_F(\nabla) + \chi_F(\tau). \quad (3)$$

Hence Ψ_F is the $F(M)$ -affine projector on $C(M)$, associated to the $F(M)$ -linear projector χ_F on $D_2^1(M)$.

Definition 1. A connection ∇ on M is called compatible with the (ap) -structure F (or is an F -connection) if it satisfies $\nabla F = 0$, where

$$(\nabla_X F)(Y) = \nabla_X F(Y) - F(\nabla_X Y)$$

for all $X, Y \in \Gamma(TM)$.

We will denote by $C_F(M)$ the set of all connections compatible with F . From the expression of $\Psi_F(\nabla)$ it follows that $\Psi_F(\nabla)_X(F) = 0$, $\forall X \in D^1(M)$, i.e. the image of any connection ∇ by the projector Ψ_F is an F -connection.

Conversely, if $\nabla_X F = 0$ it follows that $\nabla_X \circ F - F \circ \nabla_X = 0$ and so $\Psi_F(\nabla)_X = \nabla_X$, $\forall X \in D^1(M)$ i.e. $\nabla \in \text{Im } \Psi_F$.

Thus we have:

Theorem 1. The set $C_F(M)$ of the connections compatible with the (ap) -structure F is the affine submodule of $C(M)$ which is the image of the affine projector Ψ_F , i.e. $\text{Im } \Psi_F = C_F(M)$

Let ∇^0 be a fixed connection on M . Since $\text{Im } \Psi_F = C_F(M)$, then for each connection $\nabla \in C_F(M)$ there exists $\nabla' \in C(M)$ such that $\nabla' = \Psi_F(\nabla)$. But from [5], there exists $\tau \in D_2^1(M)$ such that $\nabla' = \nabla^0 + \tau$ and therefore, $\nabla = \Psi_F(\nabla^0 + \tau)$.

Then it follows:

Theorem 2. *The set $C_F(M)$ of the connections ∇ compatible with the (ap)-structure F is given by*

$$\nabla = \Psi_F(\nabla^0) + \chi_F(\tau). \quad (4)$$

where ∇^0 is a fixed connection on M and τ is an arbitrary (1, 2)-tensor field on M .

2. Main results

Let us consider F an (1, 1)-tensor field on M , such that $F^3 = F$ and $F^2 \neq I$. We will try to construct Ψ_F such that Ψ_F is a connection on M and a similar result to the *Theorem 1* holds.

For that let $G = I - 2F^2$. We have that G is also an (1, 1)-tensor field on M and

$$G^2 = (I - 2F^2)^2 = I - 4F^2 + 4F^4 = I. \quad (5)$$

So G is an (ap)-structure on M .

Hence, if we set $\Psi'_F(\nabla)_X = \frac{1}{2}(\nabla_X + G \circ \nabla_X \circ G)$ we obtain that $\Psi'_F(\nabla)_X \in C(M)$.

Definition 2. *A connection ∇ on M is called compatible with the (1, 1)-tensor field F^2 (or is F^2 -an connection) if it satisfies $\nabla F^2 = 0$.*

Theorem 3. *The set $C_{F^2}(M)$ of the connections compatible with the (1, 1)-tensor field F^2 is the affine submodule of $C(M)$ which is the image of the affine projector $\Psi'_F(\nabla)_X$, i.e. $\text{Im } \Psi'_F(\nabla) = C_{F^2}(M)$.*

Proof. First of all, we will prove that Ψ'_F takes any connection on M into an F^2 -connection. We observe that $G \circ F^2 = F^2 \circ G = -F^2$.

Let ∇ be a connection on M . We will compute

$$\Psi'_F(\nabla)_X(F^2) = \Psi'_F \circ F^2 - F^2 \circ \Psi'_F.$$

We have, consecutively,

$$\begin{aligned}
\Psi'_F \circ F^2 &= \Psi'_F (\nabla)_X F^2(Y) = \frac{1}{2} \left[\nabla_X F^2(Y) + G(\nabla_X G(F^2(Y))) \right] = \\
&= \frac{1}{2} \left[\nabla_X F^2(Y) - G(\nabla_X F^2) \right] = \frac{1}{2} \left[\nabla_X F^2(Y) - (I - 2F^2) \nabla_X F^2(Y) \right] = \\
&= \frac{1}{2} \left[\nabla_X F^2(Y) - \nabla_X F^2(Y) + 2F^2(\nabla_X F^2(Y)) \right] = F^2(\nabla_X F^2(Y)),
\end{aligned}$$

and

$$\begin{aligned}
F^2 \circ \Psi'_F &= F^2(\Psi'_F (\nabla)_X Y) = \frac{1}{2} \left[F^2(\nabla_X Y) + F^2(G(\nabla_X(Y))) \right] = \\
&= \frac{1}{2} \left[F^2(\nabla_X Y) - F^2(\nabla_X G(Y)) \right] = \frac{1}{2} \left[F^2(\nabla_X Y) - F^2(\nabla_X(Y - 2F^2(Y))) \right] = \\
&= \frac{1}{2} \left[F^2(\nabla_X Y) - F^2(\nabla_X Y) + F^2(\nabla_X F^2(Y)) \right] = F^2(\nabla_X F^2(Y)).
\end{aligned}$$

Therefore, $\Psi'_F (\nabla)_X (F^2) = 0$ and then the image of any connection ∇ by the projector Ψ'_F is an F^2 -connection.

Now, let ∇ be an F^2 -connection. Then

$$\begin{aligned}
\Psi'_F (\nabla)_X &= \frac{1}{2} (\nabla_X + G \circ \nabla_X \circ G) = \frac{1}{2} \left[\nabla_X + (I - 2F^2) \circ \nabla_X \circ (I - 2F^2) \right] = \\
&= \frac{1}{2} \left[\nabla_X + \nabla_X - 2\nabla_X \circ F^2 - 2F^2 \circ \nabla_X + 4F^2 \circ \nabla_X \circ F^2 \right] = \\
&= \frac{1}{2} \left[2\nabla_X - 4F^2 \circ \nabla_X + 4F^4 \circ \nabla_X \right] = \nabla_X.
\end{aligned}$$

Hence $C_{F^2}(M) = \text{Im } \Psi'_F$.

Let us consider F an $(1, 1)$ -tensor field on M such that $F^{2n+1} = F$ and $F^{2n} \neq I$, where n is at least 2. We will try to generalize the construction of Ψ'_F such that Ψ'_F is a connection on M and a similar result to the *Theorem 3* holds.

For that, let $H = I - 2F^{2n}$. We have that H is also an $(1, 1)$ -tensor field on M and

$$H^2 = (I - 2F^{2n})^2 = I - 4F^{2n} + 4F^{4n} = I. \quad (6)$$

So H is an (ap)-structure on M .

Thus, if we set $\Psi_F''(\nabla)_X = \frac{1}{2}(\nabla_X + H \circ \nabla_X \circ H)$ we obtain that $\Psi_F''(\nabla)_X \in C(M)$.

Definition 3. A connection ∇ on M is called compatible with the $(I, 1)$

tensor field F^{2n} (or is an F^{2n} -connection) if it satisfies $\nabla F^{2n} = 0$.

Theorem 4. The set $C_{F^{2n}}(M)$ of the connections compatible with the $(I, 1)$ -tensor field F^{2n} is the affine submodule of $C(M)$ which is the image of the affine projector $\Psi_F''(\nabla)_X$, i.e. $\Psi_F''(\nabla)_X = C_{F^{2n}}(M)$.

Proof. First of all, we will prove that Ψ_F'' takes any connection on M into an F^{2n} -connection. We observe that: $H \circ F^{2n} = F^{2n} \circ H = -F^{2n}$.

Let ∇ be a connection on M . We will compute

$$\Psi_F''(\nabla)_X(F^{2n}) = \Psi_F'' \circ F^{2n} - F^{2n} \circ \Psi_F''.$$

We have

$$\begin{aligned} \Psi_F'' \circ F^{2n} &= \Psi_F''(\nabla)_X F^{2n}(Y) = \frac{1}{2} \left[F^{2n}(\nabla_X Y) + H(\nabla_X H(F^{2n}(Y))) \right] = \\ &= \frac{1}{2} \left[F^{2n}(\nabla_X Y) - H(\nabla_X F^{2n}) \right] = \frac{1}{2} \left[F^{2n}(\nabla_X Y) - (I - 2F^{2n})\nabla_X F^{2n}(Y) \right] = \\ &= \frac{1}{2} \left[F^{2n}(\nabla_X Y) - \nabla_X F^{2n}(Y) + 2F^{2n}(\nabla_X F^{2n}(Y)) \right] = F^{2n}(\nabla_X F^{2n}(Y)). \end{aligned}$$

and

$$F^{2n} \circ \Psi_F'' = F^{2n}(\Psi_F''(\nabla)_X Y) = \frac{1}{2} \left[F^{2n}(\nabla_X Y) + F^{2n}(H(\nabla_X(Y))) \right] =$$

$$= \frac{1}{2} \left[F^{2n}(\nabla_X Y) - F^{2n}(\nabla_X Y) + F^{2n}(\nabla_X F^{2n}(Y)) \right] = F^{2n}(\nabla_X F^{2n}(Y)).$$

So $\Psi_F''(\nabla)_X(F^{2n}) = 0$ and then the image of any connection ∇ by the projector Ψ_F'' is an F^{2n} -connection.

Now, let be ∇ an F^{2n} -connection. Then

$$\begin{aligned} \Psi_F''(\nabla)_X &= \frac{1}{2}(\nabla_X + H \circ \nabla_X \circ H) = \frac{1}{2} \left[\nabla_X - (I - 2F^{2n}) \circ \nabla_X \circ (I - 2F^{2n}) \right] = \\ &= \frac{1}{2} \left[\nabla_X + \nabla_X - 2\nabla_X \circ F^{2n} - 2F^{2n} \circ \nabla_X + 4F^{2n} \circ \nabla_X \circ F^{2n} \right] = \\ &= \frac{1}{2} \left[2\nabla_X - 4F^{2n} \circ \nabla_X + 4F^{4n} \circ \nabla_X \right] = \nabla_X. \end{aligned}$$

Thus $C_{F^{2n}}(M) = \text{Im } \Psi_F''$.

Remark 1. Once we have defined an almost product structure, F , on a differentiable manifold, M , then M carries two globally complementary distributions D and D' . We denote with V and V' respectively the projection tensor fields corresponding to the two distributions D and D' . We say that X is a vector field which belongs to the distribution D if for every point p in M , we have $X_p \in D_p(M)$, where $D_p(M) = V(T_p M)$.

Let ∇ be a linear connection on the manifold M . We say that the distribution D is parallel with respect to the connection ∇ if for every vector field X which belongs to the distribution D , the vector field $\nabla_Y X$ belongs to the distribution D for every vector field Y of the manifold M .

We can define on the manifold M two linear connections by:

$$Q_X Y = V(\nabla_X V(Y)) + V'(\nabla_X V'(Y)), \quad (7)$$

$$L_X Y = V(\nabla_{V(X)} V(Y)) + V'(\nabla_{V'(X)} V'(Y)) + V[V'X, VY] + V'[VX, V'Y]. \quad (8)$$

Proposition 1. *The distributions of the almost product structure F are both parallel with respect to the connections Q and L , for every connection ∇ .*

Proof. Let Y be a vector field which belongs to the distribution D of the almost product structure F . Then we have $V'(Y)=0$ and hence $V'(Q_X Y)=0, V'(L_X Y)=0$, for every vector field X on the manifold M .

Proposition 2. *The connection Q is equal to the connection ∇ if and only if the distributions of the almost product structure F are parallel with respect to the connection ∇ .*

Proof. If the connections Q and ∇ are equal then we get that

$$V(\nabla_X V'(Y)) + V'(\nabla_X V(Y)) = 0. \quad (9)$$

By the properties of the projectors it follows that $V(\nabla_X V'(Y))=0$ and $V'(\nabla_X V(Y))=0$.

Hence, the distributions of F are parallel with respect to the connection ∇ . The converse can be verified immediately.

Theorem 5. *If one of the connections Q or L is symmetric, then the almost product structure F is integrable.*

Proof. The almost product structure F is integrable if the distributions D and D' are involutive, that is if $V'(VX, VY)=0$ and $V(V'X, V'Y)=0$ for any two vector fields X, Y on the manifold M . If the connection Q is symmetric then we have that $Q_X Y - Q_Y X = [X, Y]$.

Thus the relation $V'(VX, VY)=0$ is equivalent to the condition

$$V'Q_{YX} VY - V'Q_{VY} VX = 0. \quad (10)$$

This condition is verified by the *Proposition 1*.

Analog if L is symmetric, using the relation $V(V'X, V'Y)=0$, the result is proved

REFERENCES

1. Walker G. A., *Connections for Parallel Distribution in the Large*, Quart. J. Math. (Oxford), (2), 6: 301-208, 1955.
2. Wilmore T. J., *Parallel Distribution on Manifolds*, Proc. London. Math. Soc. (3) 6: 191-204, 1956.
3. Legrand J., *Étude d'une généralisation des structures presque-complexes sur les variétés différentiables*, Rend. Circ. Mat. Palermo, 7: 323-354, 1958.
4. Hsu, C.J., *On Some Properties of π -Structures on Differentiable Manifolds*, Tohoku Math. J. 12: 428-454, 1960.
5. Cruceanu, V., *Connections compatibles avec certaines structures sur un fibre vectoriel banachique*, Czechoslovak. Math. J. 24 (99): 126-142, 1974.
6. Yano K., *Differential Geometry on Complex and Almost Complex Spaces*, Pergamon Press, 1956.