

ON THE EXPONENTIAL DIOPHANTINE EQUATION

$$2^x + 1009^y = p^z$$

VIRGOLICI, Horia

Faculty of Mathematics and Computer Science

Spiru Haret University, Romania

hvirgolici@yahoo.com

Abstract

The aim of this paper is to give all nonnegative solutions (x, y, z) to the equation $2^x + 1009^y = p^z$, where p is a positive rational prime number with $3 \leq p \leq 997$ (we discuss 167 equations).

Keywords: *exponential Diophantine equation, Lebesgue-Nagell equation, Catalan equation*

AMS Classification: 11D61, 11Y50

1. Introduction

It is known that the equation

$$a^x + b^y = c^z \tag{1}$$

where a, b, c are prime numbers, has only finitely many solutions, but there is no algorithm to compute all the solutions (x, y, z) . Some particular cases were treated: Nagell [8] found all solutions for $\max(a, b, c) = 7$, Makowski [5], Hadano [4], Uchiyama [10], Qi Sun and Xiaoming Zhou [9], and Xiaozhuo Yang [11] determined all solutions for $11 \leq \max(a, b, c) \leq 23$. Cao [3] found all solutions for $29 \leq \max(a, b, c) \leq 97$ (60 solutions in total).

The aim of this paper is to give all nonnegative solutions to equation (1) for $a = 2, b = 1009$ (1009 representing the first prime number > 1000) and c is a rational prime number, $3 \leq c \leq 997$. The main results are given by the next theorem.

Theorem 1. *The only equations $2^x + 1009^y = p^z$, with p rational prime, $3 \leq p \leq 1009$, which admit nonnegative solutions (x, y, z) are (taking $a < b$):*

- $2^x + 1009^y = 3^z$, which has the solutions $(1, 0, 1)$ and $(3, 0, 2)$.
- $2^x + 1009^y = 5^z$, which has the solution $(2, 0, 1)$.
- $2^x + 1009^y = 17^z$, which has the solution $(4, 0, 1)$.
- $2^x + 1009^y = 257^z$, which has the solution $(8, 0, 1)$.

The proof for Theorem 1 is given in subsection 3.

2. Preliminaries

Below we present some theorems which establish the maximum number of solutions for the equation (1). Theorem 2 shows that the Catalan's equation has only one solution and gives this solution.

Theorem 2. ([1, 6, 7]). *Equation (named Catalan's equation)*

$$a^x - b^y = 1 \quad (2)$$

has no solutions in integers $a, b, x, y > 1$ other than $3^2 - 2^3 = 1$.

The theorem presented below is concerned with some particular cases of the equation (1).

Y. Bugeaud, M. Mignotte, S. Siksek solved completely the Lebesgue-Nagell equation

$$x^2 + D = y^n \quad (3)$$

where x, y are integers, $n \geq 3$ and $1 \leq D \leq 100$. From that paper, we present only the case $D = 16$ (which interests us) in the following

Theorem 3. ([2]) *If $D=16$, equation (3) has solutions $(x, y, n)=(0, 2, 4), (4, 2, 5)$. No other solutions exist in this case.*

3. Proofs of the main results

We give the proof for Theorem 1, which treats the equation

$$2^x + 1009^y = p^z, \quad 3 \leq p \leq 997, p \text{ rational prime} \quad (4)$$

Many of equations (4) have no solutions (x, y, z) due to:

Lemma 1. *If $p \equiv 1, 7, 11, 13, 19, 23, 25, 29, 31, 35, 37, 41, 43, 47 \pmod{48}$ then the equation (4) has no nonnegative solutions (x, y, z) .*

Proof: It is obvious, taking the equation (4) mod 48.

Thus, we discuss the following equations, in which p doesn't comply with the conditions in Lemma 1:

- i/ $2^x + 1009^y = p^z$, where $p \in \{53, 197, 293\}$; by taking the equation mod 24, it results that $x = 2$; by taking the equation mod 7, it results that the equation has no solutions.
- ii/ $2^x + 1009^y = p^z$, where $p \in \{101, 149\}$; by taking the equation mod 24, it results that $x = 2$; by taking the equation mod 5, it results that the equation has no solutions.
- iii/ $2^x + 1009^y = p^z$, where $p \in \{389, 401, 449, 641, 821, 881, 929\}$; by taking the equation mod 15, it results that the equation has no solutions.
- iv/ $2^x + 1009^y = 3^z$; if $y=0$ the equation has solutions $(x, y, z)=(1, 0, 1)$ (obviously) and $(3, 0, 2)$, due to Theorem 2; we consider next $y \neq 0$; by taking the equation mod 3, it results that x is odd; by taking the equation mod 16, it results one of three possibilities:

- (a) $x=1$, which leads to equation $2+1009^y = 3^z$; by taking the equation mod 9, it results that the equation has no solutions.
- (b) $x=3$, which leads to the equation $8 + 1009^y = 3^z$; by taking the equation mod 16, it results $z \equiv 2(\text{mod } 4)$; by taking the equation mod 5, it results that y is even; by taking the equation mod 13, it follows that $y \equiv 0(\text{mod } 4)$ and $z \equiv 2(\text{mod } 12)$; by taking the equation mod 17, it results that $y \equiv 0(\text{mod } 16)$ and $z \equiv 2(\text{mod } 48)$; by taking the equation mod 97, it results that $y \equiv 0(\text{mod } 96)$, and by taking the equation mod 81 it results that there are no solutions.
- (c) $z \equiv 0(\text{mod } 4)$ and $x > 4$; by taking the equation mod 5, it results that $x \equiv 1(\text{mod } 4)$ and y is odd; by taking the equation mod 7, it results that $x \equiv 0(\text{mod } 3)$ and $z \equiv 2(\text{mod } 6)$; then it results that $x \equiv 9(\text{mod } 12)$ and $z \equiv 8(\text{mod } 12)$; by taking now the equation mod 1009, it results that the equation has no solutions.

In conclusion, the equation iv/ has only the solutions (1,0,1) and (3,0,2).

- v/ $2^x + 1009^y = 5^z$; by taking the equation mod 24, it results that $x=2$ and z is odd, which leads to the equation $4 + 1009^y = 5^z$; by taking this equation mod 13, it results that $y \equiv 0(\text{mod } 4)$ and $z \equiv 1(\text{mod } 4)$; if $z=1$, the equation has solution (2,0,1); if $z \geq 3$, by taking the equation mod 7, it results that $z \equiv 1(\text{mod } 12)$; by taking the equation mod 17, it results that $y \equiv 0(\text{mod } 8)$; by taking the equation mod 25, it results that $y \equiv 8(\text{mod } 40)$; by taking the equation mod 1009, it results that there are no solutions.
- vi/ $2^x + 1009^y = 17^z$; by taking the equation mod 3, it results that x is even and z is odd; by taking the equation mod 16, it results that $x \geq 4$; by taking the equation mod 32 it results one of the possibilities:
 - (a) y is odd and $x > 4$; by taking the equation mod 17, it results that the equation has no solutions.
 - (b) $x=4$ and y is even; for $z=1$ the equation has solution (4,0,1) and for $z \geq 3$, the equation has no solutions, due to Theorem 3.
- vii/ $2^x + 1009^y = 113^z$; by taking the equation mod 7, it results that the equation has no solutions.
- viii/ $2^x + 1009^y = 257^z$; by taking the equation mod 3, it results that x is even and z is odd; by taking the equation mod 16, it results that $x \geq 4$; by taking the equation mod 32, it results one of the possibilities:
 - (a) $x=4$ and y is odd, so the equation becomes $16 + 1009^y = 257^z$; by taking this equation mod 5, it results that the equation has no solutions.
 - (b) $x \geq 6$ and y is even; if $y=0$ the equation has the solution (8,0,1) and it is the only solution in this case due to Theorem 2; if $y \geq 2$, by taking the equation mod 252, it results that $x \equiv 2(\text{mod } 6)$

- and $z \equiv 1 \pmod{6}$; by taking the equation mod 13, it results that $x \equiv 8 \pmod{12}$ and $y \equiv 0 \pmod{4}$; by taking the equation mod 1009, it results that there are no solutions.
- ix/ $2^x + 1009^y = 353^z$; by taking the equation mod 3, it results that x is even and z is odd; by taking the equation mod 16, it results that $x \geq 4$; by taking the equation mod 32, it results one of the possibilities:
- (a) $x=4$ and y is odd, so the equation becomes $16 + 1009^y = 353^z$; by taking this equation mod 5, it results that the equation has no solutions.
- (b) $x \geq 6$ and y is even; if $y=0$ the equation has no solutions; if $y \geq 2$, by taking the equation mod 5, it results that $x \equiv 0 \pmod{4}$ and $z \equiv 3 \pmod{4}$; by taking the equation mod 17, it results that there are no solutions.
- x/ $2^x + 1009^y = 593^z$; by taking the equation mod 24, it results that x is even and at least 4 and z is odd; by taking the equation mod 48, it results that y is even; if $y=0$, the equation has no solutions; if $y \geq 2$, by taking the equation mod 5, it results that $x \equiv 0 \pmod{4}$ and $z \equiv 3 \pmod{4}$; by taking the equation mod 1009, it results that there are no solutions.
- xi/ $2^x + 1009^y = 677^z$; by taking the equation mod 24, it results that $x=2$, so the equation becomes $4 + 1009^y = 677^z$, with z odd; by taking this equation mod 65, it results that the equation has no solutions.
- xii/ $2^x + 1009^y = 773^z$; by taking the equation mod 24, it results that $x=2$, so the equation becomes $4 + 1009^y = 773^z$, with z odd; by taking this equation mod 9, it results that the equation has no solutions.
- xiii/ $2^x + 1009^y = 977^z$; by taking the equation mod 3, it results that x is even and z is odd; by taking the equation mod 16, it results that $x \geq 4$; by taking the equation mod 32, it results one of the possibilities:
- (a) $x=4$ and y is even, so the equation becomes $16 + 1009^y = 977^z$, which has no solutions due to Theorem 3.
- (b) $x \geq 6$ and y is odd; by taking the equation mod 7, it results that $x \equiv 0 \pmod{3}$; by taking the equation mod 5, it results that $x \equiv 2 \pmod{4}$ and $y \equiv 1 \pmod{4}$; from these two relations, it results that $x \equiv 6 \pmod{12}$; by taking the equation mod 61 and taking into account that $x \equiv 6 \pmod{12}$ and $y \equiv 1 \pmod{4}$, it results that the equation has no solutions.

References

1. Bilu, Yu.F., *Catalan's Conjecture (after Mihăilescu)*, "Séminaire Bourbaki", Exposé 909, 55ème année, 2002-2003.
2. Bugeaud, Y., Mignotte, M., Siksek, S., *Classical and Modular Approaches to Exponential Diophantine Equations. II. The Lebesgue-Nagell Equation*, "Compos. Math.", 142, 31-62, 2006.
3. Cao, Z. F., *On the Diophantine Equation $a^x = b^y + c^z$, I*, "Chinese Sci. Bull.", 32, 1519-1521, 1987; II, *ibid.*, 33, 237, 1988 (in Chinese).
4. Hadano, T., *On the Diophantine Equation $a^x = b^y + c^z$* , "Math. J. Okayama Univ.", 19, 25-29, 1976/77.
5. Makowski, A., *On the Diophantine Equation $2^x + 11^y = 5^z$* , "Nord. Mat. Tidskr.", 7, 81-96, 1959.
6. Mihailescu, P., *A Class Number Free Criterion for Catalan's Conjecture*, "J. of Number Theory", 99, 225-231, 2003.
7. Mihailescu, P., *Primary Cyclotomic Units and a Proof of Catalan's Conjecture*, "Journal für die reine und angewandte Mathematik", 572, 167-195, 2004.
8. Nagell, T., *Sur une classe d'équations exponentielles*, "Ark. Mat.", 3, 569-582, 1958.
9. Sun, Q., Zhou, X. M., *On the Diophantine Equation $a^x = b^y + c^z$* , "Chinese Sci. Bull.", 29, 61, 1984 (in Chinese).
10. Uchiyama, S., *On the Diophantine Equation $2^x = 3^y + 13^z$* , "Math. J. Okayama Univ.", 19, 31-38, 1976/77.
11. Yang, X. Z., *On the Diophantine Equation $a^x = b^y + c^z$* , "Sichuan Daxue Xuebao", 4, 151-158, 1985 (in Chinese).

